

2014-2015 ILLINOIS SCHOOL IMMUNIZATION REQUIREMENTS

PRE 3 AND FIRST ENTRY PRE 4

Diphtheria, Pertussis, Tetanus (DTP or DTaP) – 4 doses required

- *First 3 doses received no less than 4 weeks apart

- *The interval between the 3rd and 4th dose must be at least 6 months

Polio – 3 or more doses required

- *Has received all doses at intervals of no less than 4 weeks apart

Haemophilus Influenzae Type B (Hib)

- *Has received the primary series of Hib vaccine **OR**

- *a single dose of Hib vaccine between 15 and 59 months of age.

Hepatitis B (HB) – 3 doses required

- *First two doses received no less than 4 weeks apart

- *Interval between 2nd and 3rd dose must be two months

- *Interval between 1st and 3rd dose must be at least 4 months

- *3rd dose must have been received after 6 months of age.

Varicella (chicken pox) – one dose required

- *Dose must have been received on or after 1st birthday

Measles, Mumps, Rubella (MMR) – one dose required

- *Dose must have been received on or after 1st birthday

Pneumococcal Conjugate

- *Has received the primary series of pneumococcal conjugate vaccine **OR**

- *a single dose of pneumococcal conjugate vaccine between 24 and 59 months of age

KINDERGARTEN OR 1ST GRADE (IF FIRST ENTRY INTO SCHOOL)

Diphtheria, Pertussis, Tetanus (DTP or DTaP) – 4 or more doses required

- *First 3 doses received no less than 4 weeks apart

- *Interval between the 3rd and 4th or final dose must be at least 6 months

- *Last dose must have been received on or after 4th birthday

Polio – 3 or more doses required

- *Has received all doses at intervals of no less than 4 weeks apart

- *Last dose must have been received on or after 4th birthday

Varicella (chicken pox) – 2 doses required

- *First dose must have been received on or after 1st birthday

- *Second dose received no less than 4 weeks after first dose

Measles, Mumps, Rubella (MMR) – 2 doses required

- *First dose must have been received on or after 1st birthday

- *Second dose received no less than 4 weeks after first dose

SIXTH GRADE

Diphtheria, Pertussis, Tetanus (Tdap)

- *One dose required regardless of the interval since the last DTaP, DT, or Td dose

Varicella (chicken pox) – 2 doses required

- *First dose must have been received on or after 1st birthday

- *Second dose must have been received no less than 4 weeks after first dose

Hepatitis B

- *Must show proof of 3 doses of Hepatitis B vaccine given at appropriate intervals if series had not been completed as a preschooler.